PLANTAR FIBROMA



What is a Plantar Fibroma?

A plantar fibroma is a fibrous knot (nodule) in the arch of the foot. It is embedded within the plantar fascia, a band of tissue that extends from the heel to the toes on the bottom of the foot. A plantar fibroma can develop in one or both feet, is benign (non-malignant), and usually will not go away or get smaller without treatment.

Definitive causes for this condition have not been clearly identified.


Signs and Symptoms

The characteristic sign of a plantar fibroma is a noticeable lump in the arch that feels firm to the touch. This mass can remain the same size or get larger over time, or additional fibromas may develop.

People who have a plantar fibroma may or may not have pain. When pain does occur, it is often caused by shoes pushing against the lump in the arch, although it can also arise when walking or standing barefoot.

Diagnosis

To diagnose a plantar fibroma, the foot and ankle surgeon will examine the foot and press on the affected area. Sometimes this can produce pain that extends down to the toes. An MRI or biopsy may be performed





to further evaluate the lump and aid in diagnosis.

Treatment Options


Non-surgical treatment may help relieve the pain of a plantar fibroma, although it will not make the mass disappear. The foot and ankle surgeon may select one or more of the following non-surgical options:

 Steroid injections. Injecting corticosteroid medication into the mass may help shrink it and

- thereby relieve the pain that occurs when walking. This reduction may be only temporary and the fibroma could slowly return to its original size.
- Orthotic devices. If the fibroma is stable, meaning it is not changing in size, custom orthotic devices (shoe inserts) may relieve the pain by distributing the patient's weight away from the fibroma.
- Physical therapy. The pain is sometimes treated through physical therapy methods that deliver anti-inflammatory medication into the fibroma without the need for injection.

If the mass increases in size or pain, the patient should be further evaluated. Surgical treatment to remove the fibroma is considered if the patient continues to experience pain following non-surgical approaches.

Surgical removal of a plantar fibroma may result in a flattening of the arch or development of hammertoes. Orthotic devices may be prescribed to provide support to the foot. Due to the high incidence of recurrence with this condition, continued follow-up with the foot and ankle surgeon is recommended.



This information has been prepared by the Consumer Education Committee of the American College of Foot and Ankle Surgeons, a professional society of 6,200 foot and ankle surgeons. Members of the College are Doctors of Podiatric Medicine who have received additional training through surgical residency programs.

The mission of the College is to promote superior care of foot and ankle surgical patients through education, research and the promotion of the highest professional standards.



Dr. Nigro Foot and Ankle Care Centers

The Best Available Care; Trusted by Doctors and Preferred by Patients

Main Office (Vandergrift):

127 Columbia Ave, Vandergrift, PA 15690 Phone: (724) 567-7520, Fax: (724) 568-2169

Glenshaw Office

922 Garden Place, Glenshaw, PA 15116 Phone: (412) 486-5100, Fax: (412) 486-5122

Natrona Heights Office

1601 Union Ave Suite B, Natrona Heights, PA 15065 Phone: (724) 226-0544, Fax: (724) 226-2172

Cabot Office

Concordia Professional Building 1134 Marwood Road, Cabot, PA 16023 Phone:(724) 226-0544, Fax: (724) 226-2172